

**- If you have atrial fibrillation, please contact the Nuclear Medicine Department for advice (01792 285 295) -**

**Stop 3 full days before + day of the test**

**Please follow these important instructions:**

Please bring your medication and/or repeat prescription print-out with you. You will be able to take the medication you would normally have taken on the morning of the test when you have your break at the tea-bar.

<b>Generic Name</b>	<b>Trade Name</b>
Atenolol	Tenormin, Kaltén, Tenoret 50, Tenoretic, Tenif, Co-Tenidone
Bisoprolol	Cardicor, Emtor, Monocor
Carvedilol	Eucardic
Celiprolol	Celectol
Digoxin	Lanoxin
Labetalol	Trandate
Metoprolol	Betaloc, Lopressor
Nadolol	Corgard
Nebivolol	Nebilet
Oxprenolol	Trasicor
Pindolol	Visken
Propranolol	Inderal, Inderetic, Inderex
Sotalol	Sotacor
Timolol	Betim

**Stop 2 full days before + day of the test**

<b>Generic Name</b>	<b>Trade Name</b>
Diltiazem	Tildiem, Adizem, Angitil, Calcicard, Dilcardia, Dilzem, Slozem, Viazem, Zemtard
Amlodipine	Istin
Nifedipine	Adalat, Adipine, Coracten, Fortipine, Nifedipress, Tensipine, Valri XL
Felodipine	Plendil
Isradipine	Prescal
Lacidipine	Motens
Lercanidipine	Zandip
Nimodipine	Nimotap
Verapamil	Securon, Cardilox, Univer, Verapress, Vertab
Dipyridamole	Persantin, Asasantin
Theophylline	Nuelin, Slo-phyllin, Theodur, Uniphyllin
Aminophylline	Phyllocontin
Ivabradine	Procoralan

**Stop for 12 hours before the test**

Coffee and all coffee flavoured products (including decaffeinated)
Tea (including decaffeinated, iced tea & herbal)
Chocolate and all chocolate flavoured products
Cocoa
Cola (including diet versions)
Dr Pepper
Energy Drinks e.g. Red Bull
Anadin
Panadol
Hedex
Solpadeine
Beechams
Benylin
Lemsip
Sudafed